

From: Prof G. Vidar on 22/11/99

Att: lawyer - Kubi

Fragmentation and Flow Regulation of River Systems in the Northern Third of the World

Mats Dynesius and Christer Nilsson*

Seventy-seven percent of the total water discharge of the 139 largest river systems in North America north of Mexico, in Europe, and in the republics of the former Soviet Union is strongly or moderately affected by fragmentation of the river channels by dams and by water regulation resulting from reservoir operation, interbasin diversion, and irrigation. The remaining free-flowing large river systems are relatively small and nearly all situated in the far north, as are the 59 medium-sized river systems of Norway, Sweden, Finland, and Denmark. These conditions indicate that many types of river ecosystems have been lost and that the populations of many riverine species have become highly fragmented. To improve the conservation of biodiversity and the sustainable use of biological resources, immediate action is called for to create an international preservation network of free-flowing river systems and to rehabilitate exploited rivers in areas that lack unaffected watercourses.

Natural rivers, including their riparian zones, belong to the most diverse, dynamic, and complex ecosystems on the world's continents (1). At the same time, the damming of rivers has been identified as one of the most dramatic and widespread deliberate impacts of humans on the natural environment (2). The area of former terrestrial habitat inundated by all large ($>10^8$ m³) reservoirs in the world is comparable with the area of California or France (3). Damming and diversion have greatly changed the conditions for riparian and aquatic organisms in standing as well as flowing waters (4) in three major ways: The habitats for organisms adapted to natural discharge and water-level regimes are impoverished (2, 4-7), the ability of each river to serve as a corridor is reduced (8, 9), and the function of the riparian zone as a filter between upland and aquatic systems is greatly modified (10). These adverse ecological effects have only recently been recognized (2, 5-7). The need to preserve free-flowing rivers, representing different geomorphic settings and biomes, has now been accentuated (11), and the rehabilitation of degraded rivers has been initiated in many countries (12).

Assessments of human impacts over large areas are necessary for successful environmental management, and such studies are being produced at an increasing rate (13). However, assessments of the direct human-induced changes of the river flow and river channel continuity of all large river systems are lacking (but see, for exam-

ple, (3) for gross global assessments), as are complete and reliable lists of large river systems in terms of the discharge. We compiled such data for all large river systems in the northern third of the world using a large amount of information from publications, experts, and agencies (14-17). These data were reviewed by several experts and agencies and provided the basis for sorting the river systems into three classes of exploitation: strong, moderate, and no impact. To test whether the obtained patterns also apply at a higher resolution, we made a similar study incorporating the medium-sized rivers of Norway, Sweden, Finland, and Denmark.

Background

River systems and their riparian zones play key roles in the regulation and maintenance of biodiversity in the landscapes. They have a fundamental role in the movement of organisms and dead matter (8), being the most important natural corridors through the landscape (18). Ecologists now view rivers both as systems with their own characteristics and as mediators of communication in several dimensions. Ecological interactions can be apprehended in both directions between the river and the receiving sea, the main river channel and its tributaries, the river's source and mouth (9), the river and its terrestrial surroundings (10), the river and the atmosphere, and even the river and the hyporheic water (that is, water moving underground) (19). Therefore, evaluations of the environmental impacts of human activities as well as strategies for river conservation should use the whole river basin as the basic functional unit of river landscapes (20).

The expansion of human populations and activities has resulted in extensive damming, regulation, and diversion of the world's rivers. The number of large dams in the world has increased sevenfold from 1950 to 1986, up to about 39,000 (21). The usable man-made reservoir capacity in relation to the annual river runoff is about 9% on a world basis, 10% in Europe, and 22% in North America (3). Diversion schemes have become common, and in Canada 4400 m³ of water is diverted each second and not returned to the stream of origin (22), equaling twice the discharge of the Nile. As much as 6% of the world's river runoff is evaporated through human manipulations, mainly by irrigation but also by evaporation from reservoirs (3). From the Colorado River system 64% of the runoff is consumed by irrigation, and an additional 32% is lost by evaporation from reservoirs (23); little water reaches the Gulf of California.

Size and Location

The term "river" may refer to the segment of a river channel bearing a particular name, to the entire main channel, or to a tributary. Hence, we use the term "river system" throughout to emphasize the fact that we studied entire networks of stream and river channels interconnected by surface freshwater, from the headwaters to the sea (24). There are also many different ways of defining river size. Easily assessed variables such as main channel length or catchment area are often used. However, several ecological studies suggest that the structure of riverine ecosystems is strongly correlated with water discharge (25, 26). For most rivers, the highest discharge is found close to the receiving sea, but in arid areas, discharge decreases naturally downstream. We defined a large river system (LRS) as a system that has, anywhere in its catchment, a river channel section with a virgin mean annual discharge (VMAD, the discharge before any significant direct human manipulations) of at least 350 m³ s⁻¹. We set the corresponding lower limit of a medium-sized river system (MRS) to 40 m³ s⁻¹ (27).

We limited our study to Europe, the former Soviet Union, and North America north of Mexico, or the northern third of the world's land (including Antarctica). There are 139 LRSs with either their mouths or more than half of their catchments within this area (Table 1 and Figs. 1 and 2). These LRSs have a total VMAD of nearly 254,000 m³ s⁻¹, or 20% of the world's river runoff. North America has 74 and Eurasia has 65 LRSs, but the Eurasian LRSs have a 21% higher total VMAD (139,000 versus 115,000 m³ s⁻¹) because the average Eurasian system has a 37% larger VMAD than its North American

*The authors are in the Department of Ecological Botany, Umeå University, S-901 87 Umeå, Sweden.

*To whom correspondence should be addressed.

Table 1. River systems in the northern third of the world with a VMAD of 350 m³ s⁻¹ or more (74, 75), as well as assessments of fragmentation by dams in the main channel and in tributaries and of flow regulation (76, 77) for each river system. VMAD refers to the most water-rich river channel section, in most cases close to the estuary, before any significant direct human manipulation. The main channel is the channel having the highest VMAD. The river systems are grouped by continent and impact class as defined in Table 5. The numbers beside each river system refer to those in Figs. 1 and 2. Fragmentation is ranked into five classes describing the longest main-channel segment without dams (but frequently including reservoir water tables) in relation to the entire main channel (0 = 100%; 1 = 75 to 99%; 2 = 50 to 74%; 3 = 25 to 49%; and 4 = 0 to 24%). For the tributaries, fragmentation is described by three classes (0 = no dams; 1 = dams only in the catchment of minor tributaries; and 2 = dams also in the catchment of the largest tributary). Flow regulation is described by reservoir live storage (reserv. live stor.), reservoir gross capacity (reserv. gross cap.), interbasin diversion (interb. div.), and irrigation consumption (irrig. consumpt.) for the entire river system expressed as the percent of its VMAD. The gross capacity is the total water volume that can be retained by a dam, including the bottom water that cannot be released through the lowest outlet. Live storage is the gross capacity excluding this bottom water. A "+" sign after the live storage value means that the corresponding gross capacity value represents additional reservoirs where data on live storage is lacking. Interbasin diversions are those in which water is transferred to (+) and from (-) the river system. Irrigation consumption is the water consumed by evaporation and evapotranspiration from irrigated land. Where only gross statements are available, irrigation consumption is described by "+" (minor) or "++" (major). An additional 32% of the runoff from the Colorado River and 1.3% from the Mississippi River are consumed by increased evaporation from reservoirs. With regard to the data on the Amur, Ob, Kura, Amu-Dar'ya, Ili, and Yenisey river systems, conditions in Turkey, Afghanistan, Iran, China, and Mongolia are not included in our study, but these will not change the classification for any river.

River system	VMAD (m ³ s ⁻¹)	Fragmentation		Flow regulation (% of VMAD)				River system	VMAD (m ³ s ⁻¹)	Fragmentation		Flow regulation (% of VMAD)				
		Main channel	Trib-utary	Reserv. live stor.	Reserv. gross cap.	Interb. div.	Irrig. consumpt.			Main channel	Trib-utary	Reserv. live stor.	Reserv. gross cap.	Interb. div.	Irrig. consumpt.	
<i>North America: not affected</i>																
7. Skeena	1760	0	2		0.2			4. Columbia	7500	4	2		≥48			
8. Nass	892	0	0					27. Churchill	1270	2	2	47	>125	-6		++
9. Stikine	1000	0	0					28. Nelson	2830	3	2	90	>112	-32		+
10. Taku	~600	0	0					34. Moose	1440	3	2	7	≥7			
11. Alsek	~850	0	1	2				39. Eastmain	909	1	2	12	30	-93		
12. Copper	1700	0	0					40. La Grande	1720	3	1	96	260	+97		
13. Susitna	1400	0	0					45. Koksoak	2420	2	0	51	70	-33		
14. Kvichak	590	0	0					48. Kanairiktok	~350	1	0			-37		
15. Nushagak	1000	0	0					49. Naskaupi	~350	2	0			-57		
16. Kuskokwim	1900	0	0					50. Churchill	1620	3	2	61	≥66	-20		
17. Yukon	6370	0	2		0.1			54. Manicouagan	852	3	1		≥590			
18. Kobuk	~510	0	0					56. Betsiamites	375	2	0		≥118	-2		
19. Noatak	~350	0	0					57. Saguenay	1760	3	1		≥29	-0.3		
20. Colville	~600	0	0					58. St. Lawrence	10,800	3	2		≥22	-0.8, + >1.6		1
22. Coppermine	357	0	0					59. St. John	1100	3	2	2+	1	~ -1		
23. Back	612	0	0					60. Penobscot	450	3	2		16	~ -3		
24. Thelon, Kazan	1370	0	0					61. Kennebec, Androscoggin	488	3	2		15			
25. Thaanne, Thlewiatza	507	0	0					62. Connecticut	~540	4	2		18			
26. Seal	365	0	0					68. Santee	560	3	2		60	-15		
29. Hayes	694	0	0					69. Savannah	369	3	2		96			
30. Severn	722	0	0					71. Apalachicola	~750	3	1		22			
31. Winisk	694	0	0					72. Mobile	1900	4	2		13	>0		
32. Attawapiskat	626	0	0					74. Mississippi	18,400	3	2		31	-0.2, +0.6		8
35. Harricana	473	0	0					<i>Eurasia: not affected</i>								
37. Broadback	383	0	0					84. Kalixälven,	282	0	0					
38. Rupert	878	0	0					Torneälven	373	0	1		1			
42. Povungnituk	>350	0	0					117. Uda	800	0	0					
43. Arnaud	654	0	0					118. Tavy	362	0	0					
44. R. aux Feuilles	575	0	0					119. Penzhina	720	0	0					
46. R. à la Baleine	581	0	0					120. Kamchatka	1050	0	0					
47. George	881	0	0					121. Anadyr	2020	0	0					
51. Petit Mécatina	524	0	0					123. Indigirka	1700	0	0					
52. Natashquan	422	0	0					124. Yana	970	0	0					
53. Moisie	490	0	0					126. Olenek	1090	0	0					
73. Pascagoula	~430	0	2		1.6			127. Anabar	432	0	0					
<i>North America: moderately affected</i>																
5. Skagit	475	2	1		16			128. Khatanga, Popigay	3200	0	0					
6. Fraser	3620	0	2		26			129. Taymyra	990	0	0					
21. Mackenzie	9910	1	1		≥24		-3	130. Pyasina	2260	0	0					
33. Albany	1420	1	2		≥3		-17	132. Taz	1540	0	0					
36. Nottaway	1130	0	2		≥0		-1.2	133. Pur	1050	0	0					
41. Gr. R. Baleine	665	0	1				-4	134. Nadym	610	0	0					
55. R. aux Outardes	399	1	0		≥21			136. Pechora	4100	0	0					
63. Hudson	~620	2	2		14			137. Mezen	880	0	0					
64. Delaware	~550	1	2		10		-6	139. Onega	500	0	0					
65. Susquehanna	1198	2	2		4			<i>Eurasia: moderately affected</i>								
66. Potomac	~350	3	1		3			75. Ölfusá	440	0	2	1.2	21			
67. Pee Dee	552	3	1		8			87. Narva	450	1	0	<1	3			
70. Altamaha	406	1	2		8			90. Wisla	1080	2	2	4	6			
<i>North America: strongly affected</i>																
1. Colorado	550	3	2		560		-40	64	91. Oder	580	2	2	5	≥6		
2. Sacramento, San Joaquin	1140	2	2		98		-11, +4	++	94. Rhein, Maas	2200	2	2	2+	≥2		
3. Klamath	515	3	2		31		-8	+	95. Seine	500	1	2		5		
									96. Loire	900	1	2		≥3	-0.8	0.9

Table 1 (continued)

River system	VMAD (m ³ s ⁻¹)	Fragmentation		Flow regulation (% of VMAD)				River system	VMAD (m ³ s ⁻¹)	Fragmentation		Flow regulation (% of VMAD)			
		Main channel	Tributary	Reserv. live stor.	Reserv. gross cap.	Interb. div.	Irrig. consumpt.			Main channel	Tributary	Reserv. live stor.	Reserv. gross cap.	Interb. div.	Irrig. consumpt.
<i>Eurasia moderately affected</i>								<i>Eurasia strongly affected</i>							
98, Adour	360	0	2	0.6+	0.2			92, Elbe	750	2	2	10+	5		
110, Rioni	420	2	1	<1	<1			93, Weser	360	4	2		5		
115, Amur	10,900	0	2	9	20			97, Garonne, Dordogne	1045	3	2	6+	0.9		0.3
122, Kolyma	4060	1	0	5	11			99, Duero/Douro	650	3	2	62	≅21		
125, Lena	16,900	0	1	3	7			100, Tajo/Tejo	500	4	2	50	≅34	-4	
135, Ob	12,800	2	2	9	15	0.5	<1	101, Ebro	577	4	2	46	≅9	-2	24
138, Severn, Dvina	3330	1	0	1	1			102, Rhone	1900	4	2	2+	≅7		
<i>Eurasia strongly affected</i>															
73, Thjorsá	390	3	2	13	30			103, Po	1460	3	2	4			
77, Glommavassdr.	728	4	2	16				104, Neretva	378	3	2	7	8		
78, Göta älv	554	3	2	>54				105, Drin	~350	≅3	2		≅39		
79, Dalälven	353	3	2	26				106, Danube	6450	3	2	0.1+	≅9		
80, Indalsälven	448	4	2	40				107, Dnepr	1700	≅3	≅1	35	85		9
81, Ångermanälven	481	4	2	43		-7, +7		108, Don	890	≅1	≅1	51	95		9
82, Umeälven	435	4	1	27				109, Kuban'	430	≅1	≅1	23	37		24
83, Luleälven	500	3	2	72		-2		111, Kura	850	≅3	2	44	75		35
85, Kemijoki	553	2	1	23				112, Volga	8050	4	2	34	75		2
86, Neva	2490	3	2	4+	≅26			113, Amu-Dar'ya	2200	≅1	2	14	24		50
88, Daugava	640	3	2	1	8			114, Syr-Dar'ya	1170	≅3	2	55	75		80
89, Nemunas	620	3	2	1	5			115, Ili	570	≅2	≅1	37	150		16
								131, Yenisey	20,000	≅2	2	18	58		




Fig. 1. Impact by river channel fragmentation and water flow regulation on the 14 LRSs (VMAD ≥ 350 m³ s⁻¹) of North America north of Mexico. River systems are treated as units and are represented on the map by their catchment areas. Dark areas indicate land not covered by large river systems and land

south of the study area. Diagrams present (A) total VMAD of all rivers and (B) VMAD per river system in each impact class. Impact classes are defined in Table 5, and river system numbers refer to those in Table 1. There are no LRSs on Greenland.

Table 2. River systems in Norway, Sweden, Finland, and Denmark with $40 \text{ m}^3 \text{ s}^{-1} < \text{VMAD} < 350 \text{ m}^3 \text{ s}^{-1}$ (175), and an assessment of fragmentation by dams in the main channel and in tributaries and of flow regulation (17) for each river system. The main channel is the one having the highest VMAD. River systems are grouped by impact class (Table 5). The numbers beside each system refer to Fig. 3. Fragmentation is defined in Table 1. For interbasin diversions, water transfers to (+) and from (-) the river system are indicated. There is no significant irrigation.

River system	VMAD ($\text{m}^3 \text{ s}^{-1}$)	Fragmentation		Flow regulation (% of VMAD)	
		Main channel	Tributary	Reserv. live stor.	Interb. div.
<i>Not affected</i>					
141. Tana/Tenojoki	187	0	0		
146. Saltdalsvassdraget	50	0	1	0.2	
152. Snåsavassdraget	75	0	0		
153. Verdalsvassdraget	65	0	0		+9
163. Gaularvassdraget	55	0	0		
189. Byskeälven	41	0	0		
191. Rånneälven	42	0	0		
192. Simojoki	44	0	0		
194. Kiiminkijoki	44	0	0		
<i>Moderately affected</i>					
142. Altavassdraget	84	1	0	5	
143. Reisavassdraget	43	0	1	0.7	
144. Målselvassdraget	180	0	2	21	-3.5, +0.9
147. Beiarelva	48	1	1		
150. Vefsna	161	0	1		-15
154. Stjørdalsvassdraget	82	2	1		-7
156. Gaula	98	0	1	8	
158. Surna	59	0	1	5	
159. Driva	69	0	2	21	
160. Rauma	41	0	1	13	+3
161. Breimsvassdraget	43	1	1	3	+22
162. Jølstra	54	2	0	3	
170. Bjerkreimsvassdr.	57	2	0	3	
175. Tovdalsvassdraget	64	0	2	4	
190. Piteälven	155	1	1	5	
193. Iijoki	172	2	1	4	
196. Kyrönjoki	44	2	2	9	
<i>Strongly affected</i>					
140. Pasvikelva/Paatsjoki	170	2	1	48	
145. Sulitjelmavassdraget	40	4	2	63	-13
148. Ranavassdraget	220	2	1	34	-12, +23
149. Røssåga	96	2	1	85	-4, +16
151. Namsen	270	3	1	14	-14, +13
155. Nidelvassdraget	117	3	2	38	
157. Orkla	70	3	2	19	
164. Jostedalå	60	1	1	36	
165. Årdalsvassdraget	44	1	2	26	-5
166. Vossovassdraget	104	2	1	8	-4, +15
167. Eidfjordvassdraget	50	2	1	28	-41
168. Suldalsvassdraget	103	3	2	28	-78, +68
169. Årdalselva	45	2	2	40	-55
171. Sira	116	3	2	53	-5, +44
172. Kvina	87	3	0	41	-59, +6
173. Mandalselva	85	3	2	14	
174. Otra	148	3	2	42	-6
176. Arendalsvassdraget	119	3	2	36	
177. Skjensvassdraget	287	3	2	47	
178. Numedalslågen	119	3	2	25	-0.8
179. Drammensvassdr.	326	3	2	34	+0.3
180. Åtran	47	3	2	>4	
181. Nissan	40	3	2	>0.2	
182. Lagan	72	3	2	>18	
183. Helgeån	44	3	2	>0.5	
184. Motala ström	91	3	2	>7	
185. Mälaren-Norrström	166	3	2	>5	
186. Ljusnan	219	4	2	22	
187. Ljungan	136	4	2	29	
188. Skellefteälven	155	4	2	62	
195. Oulujoki	254	3	2	52	
197. Kokemäenjoki	226	3	2	18	
198. Kymijoki	303	3	2	21	

counterpart. In Norway, Sweden, Finland, and Denmark there are 59 MRSs (Table 2, Fig. 3) contributing 38% more runoff than the nine LRSs within the same area (6500 versus $4700 \text{ m}^3 \text{ s}^{-1}$).

The catchments of the 139 LRSs span all 10 different biomes (28) of the study area, from tundra to warm desert (Table 3). The biomes are represented by 46 out of 52 different biogeographic provinces; 18 in the Nearctic and 28 in the Palearctic realm. Half of the LRSs have catchments dominated by temperate needle-leaf forests or woodlands, a biome that also covers minor parts of an additional 11% of the LRSs. The biogeographic province holding most LRSs, dominating 36 and covering parts of another 5, is the Canadian taiga. In the Palearctic realm, the West Eurasian taiga covers at least parts of 21 river systems and dominates 18 of those. The Yenisey and Ob river systems encompass the widest biogeographic variation with six biogeographic provinces each (Table 4). In North America, the catchment of the Mississippi River extends over five biogeographic provinces.

Channel Fragmentation

We assessed dam-induced fragmentation of the river channel corridor, but not, for example, fragmentation of riparian forest corridors by forestry or agriculture. All dams except low weirs were considered to have a fragmenting effect. The main channel and the tributaries were assessed separately. We measured the longest segment of the main river channel that was without dams (but that frequently included reservoir water tables) using five classes, and for tributaries we assessed the fragmentation with respect to the size of the affected tributary using three classes (see Table 1). The main channel is the channel with the highest VMAD.

At least 11 LRSs (8%) fall into the class of maximum fragmentation of both the main channel and the tributaries (29), eight of which are in Europe and three in the United States (Table 1). One additional European LRS (Umeälven) has a maximum fragmentation of the main channel but not of the tributaries. There is no fragmentation in 31 (42%) of the North American LRSs or in 18 (28%) of the Eurasian ones. Among the MRSs of the Nordic countries, four (7%) have maximum fragmentation, whereas eight (14%) are completely unfragmented (Table 2).

Flow Regulation

We assessed three types of human manipulations of the flow regime: reservoirs, inter-system water transfer, and irrigation consumption. We quantified the impact of the

reservoirs by their summed capacity, irrespective of their location in the catchment. This measure is expressed as the percentage of one average year's discharge of the river system that can be contained in the reservoirs. Available capacity data are inconsistent. Some references give "gross capacity," whereas others give "live" or "active" storage. The latter measure is preferable because it includes only the volume that can be withheld in, and subsequently released from, the reservoir. Gross capacity also includes bottom water below the invert level of the lowest outlet, so-called "dead storage," that cannot be used for regulation.

The highest live storage value is registered for La Grande Rivière in Québec (96%), and the highest gross capacity is found in Rivière Manicouagan (590%), also in Québec (Table 1). In Eurasia, the highest recorded live storage for an LRS is 72% (Luleälven in Sweden) and the highest gross capacity for an LRS is 150% (Ili in Kazakhstan). The medium-sized Røssåga system in Norway has a live storage of as much as 85% (Table 2).

We define intersystem water transfers or interbasin diversions as water transferred from one river system to another or through a man-made shortcut to the sea. Irrigation consumption refers to the amount of water evaporated or evapotranspired as a result

of irrigation and excludes water returned from irrigated land to the river system. We also express the average amount of water transferred or consumed as a percentage of one year's discharge of the river system (in

some cases only gross statements such as "major" or "minor" are available). The highest interbasin diversion percentages are also recorded for La Grande project in Québec: Rivière Eastmain loses 93% of its wa-

Table 3. Representation of biomes (28) in the catchments of the 139 largest river systems of the northern third of the world and the impact class distribution (Table 5) of the river systems that lie at least in part within each biome. For the number of river systems, the first value in parentheses represents the river systems where the biome dominates, whereas the second value in parentheses represents those where the biome covers a minor portion. Each river system flows through between one and five biomes. Biomes are listed in order of increasing degree of river system exploitation. The "northern biogeographic provinces" include the Icelandic, Subarctic birchwoods, and Kamchatkan provinces.

Biome	Number of river systems	Impact class distribution of the river systems		
		Not affected (%)	Moderately affected (%)	Strongly affected (%)
Tundra and barren arctic	42 (14 + 28)	79	10	12
Subtropical and temperate rain forests	10 (1 + 9)	70	10	20
Temperate needle-leaf forests	85 (69 + 16)	53	13	34
Temperate broad-leaf forests, and subpolar deciduous thickets				
Northern provinces	5 (3 + 2)	40	20	40
Southern provinces	37 (28 + 9)	3	38	59
Mixed mountain and highland systems	29 (16 + 13)	14	24	62
Temperate grasslands	11 (3 + 8)	0	27	73
Evergreen sclerophyllous forests	11 (4 + 7)	0	18	82
Cold-winter (continental) deserts	8 (1 + 7)	0	0	100
Lake systems	5 (0 + 5)	0	0	100
Warm deserts	1 (0 + 1)	0	0	100
All river systems		39	19	42




Fig. 2. Impact by river channel fragmentation and flow regulation on the 65 LRSs ($VMAD \geq 350 \text{ m}^3 \text{ s}^{-1}$) of Europe and the republics of the former Soviet Union. River systems are treated as units and are represented on the map by their catchments. White areas indicate land not covered by LRSs and land

south of the study area. Diagrams present (A) total VMAD of all rivers and (B) VMAD per river system in each impact class. Impact classes are defined in Table 5, and river system numbers refer to those in Table 1.

ter, and La Grande Rivière gains 97% on top of its original discharge (Table 1). In Eurasia, the highest recorded interbasin di-

version is in the Norwegian MRS Suldalsvassdraget, which gains 68% and loses 78% of its VMAD (Table 2). The largest irriga-

tion consumption (80%) is recorded for Syr-Dar'ya, causing large problems in the Aral Lake area (30), whereas cold and humid areas have little irrigation (Table 1).




Fig. 3. Impact by river channel fragmentation and flow regulation on the 59 MRSs ($40 \leq \text{VMAD} < 350 \text{ m}^3 \text{ s}^{-1}$) of Norway, Sweden, Finland, and Denmark (Denmark's largest river system has a flow of only $\sim 30 \text{ m}^3 \text{ s}^{-1}$). River systems are treated as units and are represented on the map by their catchments. White areas indicate land not covered by MRSs and land outside the study area. Diagrams present (A) total VMAD of all rivers and (B) VMAD per river system in each impact class. Unaffected tributaries $\geq 100 \text{ m}^3 \text{ s}^{-1}$ in affected large river systems are also indicated (Vindelälven in Sweden and Ounasjoki in Finland). Impact classes are defined in Table 5, and river system numbers refer to those in Table 2.

Impact Class Distribution

To summarize our data on fragmentation of the river channel and flow regulation, we classified the river systems into three levels of impact: strongly affected, moderately affected, and not affected (31). Unaffected river systems are basically those without dams in their catchments, but dams in tributaries may not be disqualifying if flow regulation is $< 2\%$ of the VMAD (Table 5). If there are dams in the main channel, the river system is never considered unaffected and if there are no dams in the main channel, it is never classified as strongly affected. All river systems with less than one-quarter of their main channel length left without dams are considered strongly affected. The limit between the strongly and moderately affected classes was chosen primarily to enhance resolution (Table 5), whereas the unaffected class was conservatively defined from an ecological point of view.

Thirty-nine percent of the LRSs, representing 23% of the total VMAD ($57,639 \text{ m}^3 \text{ s}^{-1}$), are still unaffected by river channel fragmentation and flow regulation. There are 35 LRSs, with a total VMAD of $32,710 \text{ m}^3 \text{ s}^{-1}$, left unaffected in North America, compared with 19 ($24,929 \text{ m}^3 \text{ s}^{-1}$) in Eurasia (Figs. 1 and 2). In terms of VMAD, the runoff from an average regulated and fragmented LRS is twice that of an unaffected one (2305 versus $1067 \text{ m}^3 \text{ s}^{-1}$), and the largest unaffected river system, the Yukon, is only 11th in size or about one-third the size of the Yenisey (Table 4). On the other hand, four of the seven largest river systems are only moderately affected, and two of these have dams only in their tributaries. The third largest river of the study area, the Lena in Russia, has dams

Table 4. The class of impact (Table 5) and biogeographic diversity and setting (28) of all river systems with a VMAD of $5000 \text{ m}^3 \text{ s}^{-1}$ or more (14, 15) in the northern third of the world. The numbers beside river systems refer to those shown in Figs. 1 and 2. Abbreviations: NLF, temperate needle-leaf

forests or woodlands; TG, temperate grasslands; BLF, temperate broad-leaf forests or woodlands; MMH, mixed mountain and highland systems with complex zonation.

River system	VMAD ($\text{m}^3 \text{ s}^{-1}$)	Impact	No. of biomes	Dominating biome	No. of provinces	Dominating biogeographic province
131. Yenisey	20,000	Strong	5	NLF	6	East Siberian taiga
74. Mississippi	18,400	Strong	4	TG	5	Grasslands
125. Lena	16,900	Moderate	3	NLF	4	East Siberian taiga
135. Ob	12,800	Moderate	4	NLF	6	West Eurasian taiga
116. Amur	10,900	Moderate	3	BLF	4	Manchu-Japanese mixed forest
58. St. Lawrence	10,800	Strong	3	NLF	3	Canadian taiga
21. Mackenzie	9910	Moderate	3	NLF	4	Canadian taiga
112. Volga	8050	Strong	4	NLF	5	West Eurasian taiga
4. Columbia	7500	Strong	3	MMH	4	Rocky Mountains
106. Danube	6450	Strong	3	BLF	5	Middle European forest
17. Yukon	6370	No	3	NLF	3	Yukon taiga

only in one large (Vilyui) and one very small tributary.

The unaffected LRSs, with few exceptions, flow entirely within boreal and arctic regions (Figs. 1 and 2 and Table 3). Of the 7 largest unaffected river systems, only three have dominating biomes other than taiga and tundra: the Kamchatka River of eastern Russia, and the Skeena and Stikine rivers of western Canada (Table 6). All river systems that have parts of their catchments in deserts, semideserts, or lake systems (32) are strongly affected (Table 3). Furthermore, there is no free-flowing LRS with any part of its catchment in the temperate grasslands or the evergreen sclerophyllous biomes.

Among the 59 MRSs of Norway, Sweden, Finland, and Denmark, even fewer are left free-flowing: only 15% of the MRSs, representing 9% of the VMAD (Table 2 and Fig. 3). The average unaffected MRS is half the size of the average strongly affected one (67 versus 134 $\text{m}^3 \text{s}^{-1}$). All but one of the free-flowing MRSs have a VMAD of as little as 75 $\text{m}^3 \text{s}^{-1}$ or less. The Nordic MRSs show the same latitudinal gradient of exploitation (Fig. 3) as do the LRSs on the global scale. Some river systems in the north have remained unaffected, whereas southern river systems, especially in the temperate, broad-leaf biome, are strongly affected. Only two out of 18 river systems with VMADs larger than 40 $\text{m}^3 \text{s}^{-1}$ within this biome are moderately affected, namely, Tovdalsvassdraget and Bjerkreimsvassdraget, both in Norway, and none is free-flowing. Among the fragmented and regulated river systems, we found only two unaffected tributaries larger than 100 $\text{m}^3 \text{s}^{-1}$ (Fig. 3), both located in the north: Vindelälven (200 $\text{m}^3 \text{s}^{-1}$) in Sweden and Ounasjoki (140 $\text{m}^3 \text{s}^{-1}$) in Finland.

Implications for Conservation Management

Large areas in the northern third of the world completely lack unregulated LRSs. Although river exploitation may have different effects in different rivers, some inevitable consequences stand out. For example, several types of important habitats, such as waterfalls, rapids, and floodplain wetlands, may disappear from entire regions. The loss of waterfalls and rapids indicates the loss of numerous species of plants and animals specific to running waters. Wetland losses are especially serious in dry areas where alternative habitats are scarce. As a result of habitat destruction and obstruction to organism dispersal, many riverine species may have become extinct over vast areas, whereas populations of others have become fragmented and run the risk of future extinctions. Although regional depletion of river

faunas has been demonstrated and habitat fragmentation by dams has been shown to be a major cause of this depletion (33), there is insufficient knowledge of the general stage of this depletion. The final result of the present exploitation is also difficult to predict because the time needed for re-equilibration of ecosystems is probably longer than the time river regulation has been in practice (2, 34).

Would a higher resolution of river conditions change this conclusion? For example, it is not certain that an LRS classified as strongly impacted is so in every part of its catchment. However, low contribution of reasonably large, unaffected tributaries in affected river systems in Norway, Sweden, Finland, and Denmark suggests that such remaining parts are generally small. The largest strongly affected river systems in the

study (Table 4) might have large, unaffected tributaries or parts of tributaries. However, these are not complete rivers because they have no direct connections with the sea; they therefore lack, for example, estuarine flora and fauna.

River systems smaller than those we considered are also unlikely to match the losses in regional ecosystem diversity that have befallen the LRSs. Small rivers have a species composition different from that of large rivers (25, 35), and as suggested by the study of Norway, Sweden, Finland, and Denmark, MRSs may show the same geographic pattern of human impact as do LRSs.

To improve the conservation of biodiversity and the sustainable use of biological resources, immediate action is required (36). River conservation efforts should be based on the main tenet of the World Con-

Table 5. Principles for constructing three classes of river system exploitation (not affected, moderately affected, and strongly affected) from the combination of fragmentation and flow regulation assessments. The fragmentation classes are defined in Table 1. Summed values of reservoir live storages, interbasin diversions (irrespective of direction) and irrigation consumption are given as the percentage of VMAD (Table 1). If data on live storage are lacking, half the gross capacity is used as a substitute (Table 1).

Fragmentation (Main channel + tributaries)	Flow regulation (%)		
	Not affected	Moderately affected	Strongly affected
0 + 0	0		
0 + 1	≤2	>2	
0 + 2	≤1	>1	
1 + 0		≤30	>30
1 + 1		≤25	>25
1 + 2, 2 + 0		≤20	>20
2 + 1		≤15	>15
2 + 2, 3 + 0		≤10	>10
3 + 1		≤5	>5
3 + 2, 4 + 0, 1, 2			≥0

Table 6. The biogeographic diversity and setting (28) of all free-flowing river systems with a VMAD of 1000 $\text{m}^3 \text{s}^{-1}$ or more in the northern third of the world. The numbers by the river systems refer to those in Figs. 1 and 2. Abbreviations: NLF, temperate needle-leaf forests or woodlands; TUN, tundra communities and barren arctic desert; BLFn, northern temperate broad-leaf forests or woodlands as described in Table 3; MMH, mixed mountain and highland systems with complex zonation.

River system	VMAD ($\text{m}^3 \text{s}^{-1}$)	No. of biomes	Dominating biome	No. of provinces	Dominating biogeographic province
17. Yukon	6370	3	NLF	3	Yukon taiga
136. Pechora	4100	2	NLF	2	West Eurasian taiga
128. Khatanga	3200	2	NLF	2	East Siberian taiga
130. Pyasina	2260	2	TUN	3	Low-Arctic tundra
121. Anadyr	2020	2	TUN	3	Low-Arctic tundra
16. Kuskokwim	1900	2	NLF	2	Yukon taiga
7. Skeena	1760	2	MMH	2	Rocky Mountains
12. Copper	1700	2	NLF	2	Yukon taiga
123. Indigirka	1700	2	NLF	3	East Siberian taiga
9. Stikine	1600	2	MMH	2	Rocky Mountains
132. Taz	1540	1	NLF	1	West Eurasian taiga
13. Susitna	1400	1	NLF	1	Yukon taiga
24. Thelon, Kazan	1370	2	TUN	2	Canadian tundra
126. Olenek	1090	2	NLF	3	East Siberian taiga
120. Kamchatka	1050	1	BLFn	1	Kamchatkan
133. Pur	1050	2	NLF	2	West Eurasian taiga
15. Nushagak	1000	2	TUN	2	Alaskan tundra

ervation Strategy: the maintenance of ecological processes (11). Water flow and water-level fluctuation are most important because various other processes depend on them (8, 37). To maintain a near-natural water transport through the landscape, we should not limit our scope to the river channel but should include the entire catchment (38). For example, the ability of the catchment to dampen large natural fluctuations in rainfall and snowmelt must not be reduced. Today, naturally regulated runoff patterns in the catchment are being indirectly deregulated, especially by embankments, draining, and deforestation but also by urbanization and industry (39).

Different challenges are faced in different biomes. In the northern world's tundra and taiga biomes as well as in the temperate rain forest areas of Alaska and Canada, it is still possible to create an international preservation network of representative, unregulated, and unfragmented LRSs. Unless an entire catchment can be protected in one large reserve, a network of reserves is necessary to protect its entire range of biotic diversity (40). In the other biomes, the magnitude of impact indicates that river channel fragmentation and water regulation have had profound ecological effects on the LRSs (2, 5-7). Here, legislation forbidding dams on remaining free-flowing tributaries and main-channel reaches is a minor tool. More important is to find ways to minimize the negative effects of existing dams and diversions and also to consider the pervasive influences of pollution, riparian logging and agriculture, poor land-use practices within the catchment, and the invasion or introduction of nonnative biota (41). All of these influences fragment and otherwise compromise natural attributes and processes in river ecosystems and add to the effects of dams and diversions. Fully integrated catchment management, including measures of ecological rehabilitation, is needed (20, 42, 43).

Ecological rehabilitation of degraded river systems may include a variety of measures. For example, the rehabilitation of channel sinuosity will increase the retention of matter and energy (44). The rehabilitation of naturally functioning floodplains is essential to provide wildlife habitat and help reduce or buffer non-point-source pollution (45). This change necessitates both the reconnection of the river with its floodplains and the modification of water-level fluctuations toward more natural conditions (46). Examples of ongoing projects exist along the Danube and Rhine rivers in Germany and along the Mississippi River in the United States (47). The rehabilitation of migration routes and spawning sites is essential to recover anadromous salmonids; such work is under way in the Columbia

River, for example (42). In some cases, the removal of dams is also considered, as for the Elwha River of the Olympic Peninsula in the state of Washington, United States, where authorities are considering tearing down two dams, 30 and 70 m high (48). River rehabilitation measures must be based on a synthesis of ecological principles and include a basic research component to allow an adaptive refinement of management objectives as new information becomes available (49).

So far, river conservation management has mainly been a national concern. For example, legislations and plans to preserve rivers have been introduced in the United States, Canada, Norway, Sweden, and Finland (50). In this way, the main channels of the Seal, Thelon, and Kazan, the major part of the Alsek and Noatak rivers (all in North America), and the entire river system of Torne-Kalix (in Fennoscandia) have been protected. In addition, many minor parts of other LRSs are protected, and in some cases national borders also form boundaries of protection, as for the Alsek River of Alaska and western Canada (protected only in Canada) and for Vefsna, an MRS shared by Sweden and Norway (protected only in Sweden). Time has now come to adopt an international approach to river conservation and manage entire river systems irrespective of political borders.

REFERENCES AND NOTES

- R. J. Naiman, H. Décamps, M. Pollock, *Ecol. Appl.* 3, 209 (1993).
- G. E. Petts, *Impounded Rivers: Perspectives for Ecological Management* (Wiley, Chichester, UK, 1984).
- M. I. L'vovich et al., in *The Earth as Transformed by Human Action*, B. L. Turner II et al., Eds. (Cambridge Univ. Press, Cambridge, 1990), chap. 14.
- M. B. Bain, J. T. Finn, H. E. Boothe, *Ecology* 69, 382 (1988); J. E. Brittain and S. J. Saltveit, *Regul. Rivers* 3, 191 (1989); C. Nilsson and M. Dynesius, *ibid.* 9, 45 (1994); C. Nilsson, A. Ekblad, M. Gardfjell, B. Carlberg, *J. Appl. Ecol.* 28, 963 (1991); B. Rorslett, *Hydrobiologia* 175, 65 (1989).
- R. M. Baxter and P. Glaude, *Can. Bull. Fish. Aquat. Sci.* 205, 1 (1980); J. F. Craig and J. B. Kemper, Eds., *Regulated Streams: Advances in Ecology* (Plenum, New York, 1987).
- A. Lillehammer and S. J. Saltveit, Eds., *Regulated Rivers* (Universitetsforlaget, Oslo, 1984).
- G. E. Petts, P. Armitage, A. Gustard, Eds., *Regul. Rivers* 3, 1 (1989); J. A. Stanford and F. R. Hauer, Eds., *ibid.* 8, 1 (1993); J. V. Ward and J. A. Stanford, *The Ecology of Regulated Streams* (Plenum, New York, 1979).
- G. P. Malanson, *Riparian Landscapes* (Cambridge Univ. Press, Cambridge, 1993).
- J. V. Ward and J. A. Stanford, in *Dynamics of Lotic Ecosystems*, T. D. Fontaine and S. M. Bartell, Eds. (Ann Arbor Science, Ann Arbor, MI, 1983), pp. 29-42; J. V. Ward and J. A. Stanford, in *Regulated Streams: Advances in Ecology* (Plenum, New York, 1987), pp. 391-409.
- H. Décamps, *Ecol. Appl.* 3, 441 (1993).
- J. Lubchenko et al., *Ecology* 72, 371 (1991); R. L. Naiman et al., Eds., *The Freshwater Imperative: A Research Agenda* (Island Press, Washington, DC, in press); *World Conservation Strategy* (International Union for Conservation of Nature and Natural Resources, United Nations Environment Programme (UNEP), and the World Wildlife Fund (WWF), Gland, Switzerland, 1980).
- P. J. Boon, P. Calow, G. E. Petts, Eds., *River Conservation and Management* (Wiley, Chichester, UK, 1992).
- For example, L. Hannah, D. Lohse, C. Hutchinson, J. L. Carr, A. Lankerani, *Ambio* 23, 246 (1994); J. M. McCloskey and H. Spalding, *ibid.* 18, 221 (1989); D. Skole and C. Tucker, *Science* 260, 1905 (1993).
- Data on VMAD in North America are compiled from W. P. Bartlett Jr., W. B. Higgins, R. G. Lippert, *U.S. Geol. Surv. Water Data Rep. ME-91-1* (1991); *Discharge of Selected Rivers of the World*, vols. 1-3 of *Studies and Reports on Hydrology*, 5 [United Nations Educational, Scientific and Cultural Organization (UNESCO), Paris, 1969-1985]; *The Hydrological Atlas of Canada* (Fisheries and Environment Canada, Ottawa, 1978), plate 22; D. W. Moody, E. B. Chase, D. A. Aronson, *U.S. Geol. Surv. Water-Supply Pap.* 2300 (1986); P. R. Saebel, F. P. Kapinos, G. L. Knapp, *ibid.*, 2294 (1987); and C. W. Stockton and W. R. Boggess, *Transportation Research Record* 922 (U.S. National Research Council, Washington, DC, 1983). We also compiled data supplied by Environment Canada, Ottawa; Ministère de l'Environnement, Sainte-Foy, Québec; U.S. Geological Survey, Anchorage, AK; U.S. Geological Survey, Columbia, SC.
- Data on VMAD in Eurasia are compiled from "Avlepsforhold i Norske Vassdrag", *VH-Notat 6/88* (National Freshwater and Energy Authority, Oslo, Norway, 1988); J. Cottillon, A. Bartle, J. A. Veltrop, Eds., *Dams in Europe & USSR: A Geographical Approach* (International Commission on Large Dams, Paris, and International Water Power and Dam Construction, Sutton, UK, 1990); *Discharge of Selected Rivers of the World*, in (14); *Eau 2000 Réussir Votre Futur: Bassin de la Dordogne* (Agence de l'Eau Adour Garonne, Toulouse, France, 1991), booklet; *Great Soviet Encyclopedia* (Macmillan, New York, translation of Russian ed. 3, 1973-1983); *Hydrological Yearbook 1978-1989* (National Board of Waters and the Environment, Helsinki, 1981-1992); "Opplysninger om Nord-Norske Vassdrag med Middelavløp >40 m³/s," *VRN-H-Notat 4/1993* (National Freshwater and Energy Authority, Northern Region, Narvik, Norway, 1993); *Surface Water Resources of the USSR* (Gidrometeoizdat, Leningrad, 1966-1973), vols. 1-20; *Streamflow Records of Sweden* (Swedish Meteorological and Hydrological Institute, Stockholm, 1979); *Water Resources of the USSR and Their Use* (Gidrometeoizdat, Leningrad, 1987); and *Yearbook of Environmental Statistics 1986-1987* (Statistics Sweden, Stockholm, 1987). We also compiled data supplied by J. V. P. Hrvatska Vodoprivreda Zagreb OJ Split, Split, Croatia; Det Danske Hedeselskab, Slagelse, Denmark; Centre d'Ecologie des Ressources Renouvelables, Toulouse, France; Bundesanstalt für Gewässerkunde, Koblenz, Germany; National Power Company, Reykjavik, Iceland; National Freshwater and Energy Authority, Oslo, Norway; Institute of Meteorology and Water Management, Warsaw; Confederación Hidrográfica del Ebro, Zaragoza, Spain; and Swedish Meteorological and Hydrological Institute, Norrköping.
- Data on river channel fragmentation and water regulation in North America were compiled from (22); P. P. Harper, in *The Rivers Handbook*, P. Calow and G. E. Petts, Eds. (Blackwell, Oxford, 1992), vol. 1, chap. 19; *Hydraulic Schematic Diagram of the Nelson and Churchill Drainage Basins* (Manitoba Hydro, System Operative Department, Winnipeg, Canada, 1986), map; *Hydrological Atlas of Canada*, in (14), plate 34; P. P. Micklin, in *Large-Scale Water Transfers: Emerging Environmental and Social Experiences*, G. N. Gobulev and A. K. Biswas, Eds. (UNEP Water Resources Series, Tycooly Publishing, Oxford, 1985), vol. 7, pp. 37-65; *National Atlas of Canada* (Energy, Mines and Resources Canada, Ottawa, ed. 5, 1985), map 5.1; D. W. Moody, E. B. Chase, D. A. Aronson, in (14); *Operating Hydro-Electric Power Stations in Ontario* (Ministry of Energy, Ontario, Canada, 1986), map; *Overview of the Coppermine River Basin* (Environment Canada, Yellowknife, 1988); *River Basins of the United States* (U.S. Geological Sur-

- vey, Denver, CO, 1981-1985; five booklets covering the Colorado, Columbia, Delaware, Hudson, and Potomac river basins; D. Roy et al., *Ecological Monitoring Program of the La Grande Complex 1978-1984: Summary Report* (Société d'Énergie de la Baie James, Montreal, Canada, 1986); B. C. Ruddy and K. J. Hitt, *U.S. Geol. Surv. Open-File Rep.* 90-163 (1990); K. J. Hitt, *Surface Water and Related Land Resources Development in the United States and Puerto Rico* (U.S. Geological Survey, Reston, VA, 1984), map; *Surface Water Data Reference Index Canada 1983* (Environment Canada, Ottawa, 1989); *World Register of Dams* (International Commission on Large Dams, Paris, 1973-1988); and P. Yee and J. Lloyd, *Great Lakes Water Levels* (Environment Canada, Ottawa, 1989; third revised printing), booklet. We also compiled data supplied by D. Rosenberg, Winnipeg, Canada; British Columbia Hydro and Power Authority, Vancouver, Canada; Environment Canada, Winnipeg; Environment Canada, New Brunswick; Ontario Hydro, North York and Toronto, Canada; Hydro Québec and Ministère de l'Environnement, Sainte-Foy, Canada; Water Survey of Canada, Whitehorse; U.S. Geological Survey, Anchorage, AK; and U.S. Geological Survey, Office of the Delaware River Master, Milford, PA.
17. Data on river channel fragmentation and water flow regulation in Eurasia are compiled from *Bundeswasserstrassen: Informationen für die Sportsschifffahrt* (Wasser- und Schifffahrtsverwaltung des Bundes, Bonn, 1993), map; J. Cotillon, A. Bartle, J. A. Veltrop, in (15); H. Décamps, in (6), p. 497; *Eau 2000 Réussir Votre Futur*, in (15); four booklets covering the Dordogne, Adour, Tarn-Aveyron, and Lot basins; anonymous, *Tech. Sci. Munic.* 12, 651 (1993); *La Garonne Encore Sauvage, Vivante Pour Tous!* (Agence de l'Eau Adour Garonne, Toulouse, France, 1989), booklet; D. G. de Jalou, in (6), p. 482; *Mapa de Embalses* (Direccion General de Obras Hidraulicas, Madrid, 1986), map; G. Arpi, in *National Atlas of Sweden* (Swedish Society for Anthropology and Geography, Stockholm, 1966), maps 39A and 40A; "Om Samlet Plan for Vassdrag," *St. Meld. Nr. 60/1991-92* (Ministry of Environmental Protection, Oslo, Norway, 1992); "Oplysninger om Nord-Norske Vassdrag med Middelavlop >40 m³/s," *VFN-H-Nytt* 47:993 (National Freshwater and Energy Authority, Northern Region, Narvik, Norway, 1993); anonymous *Wasser Energie Luft-Eau Énergie Air* 75, 16 (1983); C. Bernes and C. Grunsten, Eds., *Sveriges Nationalatlas: Miljön* (SNA Förlag, Stockholm, 1991), pp. 114-121; "Utbygd Vannkraft pr. 01.01.1990," *NVE Publikasjon nr V 32* (National Freshwater and Energy Authority, Oslo, Norway, 1990); "Vattenkraft och Miljö 3," *SOU 1976:28* (Ministry of Housing and Physical Planning, Stockholm, Sweden, 1976); and *World Register of Dams*, in (16). We also compiled data supplied by the Public Enterprise for Generation, Transmission and Distribution of Electric Power in Bosnia and Herzegovina, Sarajevo, Bosnia; J. V. P. urvatska Vodoprivreda Zagreb OJ Split, Split, Croatia; Water Research Institute, Prague, Czech Republic; Finnish Committee on Large Dams, Vantaa; Éric Tabacchi, Toulouse, France; Établissement Public D'Aménagement de la Loire et de ses Affluents, Orléans, France; Bundesanstalt für Gewässerkunde, Koblenz, Germany; Institut für Wasserbau, Universität Stuttgart, Germany; National Energy Authority, Reykjavik, Iceland; National Power Company, Reykjavik, Iceland; Po River Basin Authority, Parma, Italy; National Freshwater and Energy Authority, Oslo, Norway; Consulting Engineers Hydroproject, Warsaw; Institute of Meteorology and Water Management, Warsaw; State Hydrological Institute, St. Petersburg, Russia; Confederacion Hidrographica del Ebro, Zaragoza, Spain; Spanish Committee on Large Dams, Madrid; Dalälvens Vattenregleringsförelägg, Borlänge, Sweden; Helgeåns Vattenvårdsförbund, Kristianstad, Sweden; Skellefteåns Vattenregleringsföretag, Skellefteå, Sweden; Urneälvens Angermanälvens, Indalsälvens, Ljungans och Lusnans Vattenregleringsföretag, Vällingby, Sweden; Vattenfall Norrbotten, Luleå, Sweden.
 - R. T. Forman and M. Godron, *Landscape Ecology* (Wiley, New York, 1986).
 - J. A. Stanford and J. V. Ward, *J. N. Am. Benthol. Soc.* 12, 48 (1993); J. V. Ward, *ibid.* 8, 2 (1989).
 - F. B. Löttspeich, *Water Resour. Bull.* 16, 581 (1980); Å. Sundborg, in (6), pp. 455-466; J. A. Stanford and J. V. Ward, in *Watershed Management*, R. J. Naiman, Ed. (Springer-Verlag, New York, 1992), chap. 5.
 - World Register of Dams* (International Commission on Large Dams, Paris, 1988; 1988 updating).
 - J. C. Day and F. Quinn, in *Proceedings of the Symposium on Interbasin Transfer of Water: Impacts and Research Needs for Canada* (National Hydrology Research Institute, Saskatoon, Canada, 1987), pp. 43-56.
 - D. W. Moody, E. B. Chase, D. A. Aronson, in (14), p. 63.
 - The definition indicates that two rivers with significant connections of natural surface freshwater through river-mouth deltas or through upstream natural cross-channels, so-called bifurcations, comprise one river system. This definition is ecologically reasonable because freshwater organisms can migrate between such rivers. Examples are the Torne-Kalix river system in Sweden and Finland, the Rhine-Maas river system in northwestern continental Europe, and the Sacramento-San Joaquin river system of California. Tributaries are not regarded as complete river systems because they have no connection to the sea.
 - G. W. Minshall, R. C. Petersen Jr., C. F. Nimz, *Am. Nat.* 125, 16 (1985).
 - R. J. Naiman, H. Décamps, J. Pastor, C. A. Johnston, *J. N. Am. Benthol. Soc.* 7, 289 (1988); B. Stutzner and B. Higl, *Freshwater Biol.* 16, 127 (1986); R. L. Vannote, G. W. Minshall, K. W. Cummins, J. R. Sedell, C. E. Cushing, *Can. J. Fish. Aquat. Sci.* 37, 130 (1980).
 - We used a global runoff map [F. van der Leeden, Ed., *Water Resources of the World—Selected Statistics* (Water Information Center, Port Washington, NY, 1975), figure 7:4] to determine, for different regions, the minimum catchment area that is required to produce an LRS. We then sorted out LRS candidates with sufficient catchment areas using operational navigation charts (1:1,000,000) of the U.S. Department of Defense Mapping Agency, a map series covering the entire world on the same scale. We checked the candidates by collecting a large amount of streamflow information from publications and then let experts and agencies review the list (14, 15). The same kind of procedure was used to prepare a list of the MRSs in Sweden, Norway, Finland, and Denmark.
 - Biomes and biogeographical provinces according to M. D. F. Udvardy, *IUCN Occasional Paper Number 18* (Morges, Switzerland, 1975).
 - In a river system with maximum fragmentation, the longest main channel segment without dams is <25% of the entire main channel length and there is at least one dam in the catchment of the largest tributary.
 - P. P. Micklin, *Science* 241, 1170 (1988).
 - We merged the assessed variables in two steps. In the first step, we integrated the fragmentation and flow regulation variables into one figure each. We combined the two fragmentation variables (main channel and tributaries) into 15 classes from 0 (no fragmentation) to 4 + 2 (maximum fragmentation). We summarized the flow regulation by adding the reservoir live storage, intersystem diversion, and irrigation consumption values into one percentage value. If water was transferred both into and out of the river system, the percentages were summed, irrespective of the direction of transfer (for example, +5% and -15% makes 20%). The ratio between live and dead storage varies among reservoirs where both are known, but in most cases, this ratio is considerably below 1. Hence, for river systems where only gross capacity is known, we halved the percentages to get an approximate live storage value to use in the classification. The second step was the amalgamation of fragmentation and flow regulation into one of the three classes of impact (Table 5).
 - Lake systems in the study area that are large and ancient lakes that may warrant special consideration as biogeographic provinces were considered, including the Great Lakes in the United States, Lake Ladoga in Russia, Aral Sea in Kazakhstan, and Lake Baikal in Russia (28).
 - V. J. Birstein, *Conserv. Biol.* 7, 773 (1993); B. R. Davies and K. F. Walker, Eds., *The Ecology of River Systems* (Junk, Dordrecht, the Netherlands, 1986); C. A. Frissell, *Conserv. Biol.* 7, 342 (1993); M. Peñáz, F. Kubiček, P. Marvan, V. Zárka, *Acta Sci. Nat. Acad. Sci. Bohemoslovaca Brno* 2, 1 (1968); M. Peñáz, J.-M. Olivier, G. Carré, D. Pont, A.-L. Roux, *ibid.* 25, 1 (1991); A. L. Roux, *Bull. Ecol.* 7, 475 (1976); A. L. Sheldon, *Conserv. Biol.* 2, 149 (1988); J. D. Williams, M. L. Warren Jr., K. S. Cummings, J. L. Harris, R. J. Neves, *Fisheries* 18, 6 (1993).
 - A. M. Milner, in *The Rivers Handbook*, P. Calow and G. E. Petts, Eds. (Blackwell, Oxford, 1994), vol. 2, chap. 5.
 - C. Nilsson et al., *J. Ecol.* 82, 261 (1994).
 - N. A. Robinson, Ed., *Agenda 21: Earth's Action Plan* (Oceana Publications, New York, 1993).
 - C. Nilsson et al. in *Current Topics in Botanical Research*, J. Menon, Ed. (Council of Scientific Research Integration, Trivandrum, India, in press).
 - R. J. Naiman, Ed., *Watershed Management* (Springer-Verlag, New York, 1992).
 - J. F. Franklin, *ibid.*, chap. 3; E. L. Turner et al., Eds., *The Earth as Transformed by Human Action* (Cambridge Univ. Press, Cambridge, 1990).
 - C. Margules, Ed., *Biol. Conserv.* 50, 1-303 (special issue); C. Nilsson, in *Ecological Principles of Nature Conservation*, L. Hansson, Ed. (Elsevier, London, 1992), chap. 9.
 - J. D. Allan and A. S. Flecker, *BioScience* 43, 32 (1993).
 - 1994 Columbia River Basin Fish and Wildlife Program Overview* (Northwest Power Planning Council, Portland, OR, 1994).
 - B. Doppelt, M. Scurlock, C. Frissell, J. Karr, *Entering the Watershed* (Island Press, Washington, DC, 1993).
 - G. Pinay, H. Décamps, E. Chauvet, E. Fustec, in *The Ecology and Management of Aquatic-Terrestrial Ecosystems*, R. J. Naiman and H. Décamps, Eds. (UNESCO, Paris, 1990).
 - A. R. G. Large and G. E. Petts, in (34), chap. 21; R. C. Petersen Jr, et al., *Ambio* 16, 166 (1997).
 - P. B. Bayley, *Regul. Rivers* 6, 75 (1991); G. E. Petts and I. Maddock, in (34), chap. 15.
 - K. Kern, in *Lowland Floodplain Rivers*, P. A. Catling and G. E. Petts, Eds. (Wiley, Chichester, UK, 1992), chap. 13; P. Larsen, in (34), chap. 22; A. Robinson and R. Marks, *Restoring the Big River: A Clean Water Act Blueprint for the Mississippi* (Izaak Walton League of America and Natural Resources Defense Council, Minneapolis, MN, 1994).
 - L. Sklar, *World Rivers Rev.* 8, 9 (1993).
 - C. S. Holling, *Ecol. Appl.* 3, 352 (1993).
 - The National Wild and Scenic Rivers System (United States), the Canadian Heritage Rivers System, the Norwegian Protection Plan for River Systems (Verneplan for vassdrag), the Natural Resources Act (Naturresurslagen, Sweden), and the Rapids Protection Act (Forsskyddslagen, Finland).
 - Supported by WWF in Sweden and the Swedish Society for the Conservation of Nature. We thank all the people who directed us to the experts. We acknowledge the following persons and agencies for supplying and reviewing data: Ø. Ålbu and R. Dahl (Direktoratet for Naturforvaltning, Trondheim, Norway); G. Barabé and M. Mirón (Direction du Milieu Hydrique, Sainte-Foy, Canada); R. D. Barnes (Newfoundland and Labrador Hydro, St. John's, Canada); M. Balteazzore (Istituto de Ricerca Sulle Acque, Brugherio, Italy); E. Bičakčić (Public Enterprise for Generation, Transmission, and Distribution of Electric Power in Bosnia and Herzegovina, Sarajevo, Bosnia); B. Bigelow and R. D. Lamke (U.S. Geological Survey, Anchorage, AK); R. Bischof (Swiss National Committee on Large Dams, Zürich); V. Bogatov (Far Eastern Scientific Center, Vladivostok, Russia); C. W. Boning, W. Kirby, E. N. Pickering (U. S. Geological Survey, Reston, VA); F. Bultot (Royal Meteorological Institute of Belgium, Bruxelles); Bundesanstalt für Gewässerkunde (Koblenz, Germany); Bundesministerium für Land- und Forstwirtschaft (Wien, Austria); J. M. Cucc (University of Calgary,

Canada); J. Debelius (German Committee on Large Dams, Düsseldorf); H. Décamps, F. Gazelle, G. Pinay, and E. Tabacchi (Centre d'Écologie des Systèmes Fluviaux, Toulouse, France); D. G. de Jalon (Universidad Politécnica de Madrid, Spain); F. T. Direito (Ministério do Ambiente e dos Recursos Naturais, Lisboa, Portugal); I. C. Dirom and P. Eng (British Columbia Hydro and Power Authority, Vancouver, Canada); J. E. Eaton (Ontario Hydro, North York, Canada); A. A. Elizalde (Confederacion Hidrografica del Ebro, Zaragoza, Spain); C. M. Escartin (Centro de Estudios Hidrográficos, Madrid, Spain); M. V. Getsen (Biological Institute, Syktyvkar, Russia); J. Giesecke (Universität Stuttgart, Germany); H. Goertz (Environment Canada, Guelph); M. Gotthardsson, A. Jansson, and T. Lindkvist (Swedish Meteorological and Hydrological Institute, Norrköping); M. E. Greco (Canadian Heritage Rivers Secretariat, Hull); R. Gregory (Water Survey of Canada, Whitehorse); A. Gustard (Institute of Hydrology, Wallingford, UK); M. Hare (Ontario Hydro, Toronto, Canada); W. E. Harkness (U.S. Geological Survey, Millford, PA); K. D. Harvey, A. R. Perks, and J. Power (Environment Canada, Ottawa); K. Hauge and I. Iversen (Det Kongelige Miljøverndepartement, Oslo, Norway); International Centre of Water Studies (Amsterdam, the Netherlands); J. Irvine (Pacific Biological Station, Nanaimo, Canada); A. Jacewicz and R. Ramm (Consulting Engineers Hydroproject, Warsaw); H. Jónatansson, G. Júlíusson, and A. Olsen (National Power Company, Reykjavik, Iceland); Z. Kajak (Polish

Academy of Sciences, Lomanski); E. Kielland, S. Krog, R. Skretteberg, A. Tolla, A. M. Tvede, B. Wold, and K. Wold (Norges Vassdrags og Energiwerk, Oslo, Norway); R. Klejnowski (Institute of Meteorology and Water Management, Warsaw); M. Kowalchuk (Environment Canada, Winnipeg); W. L. Kreuder (Environment Canada, Vancouver); R. Kwartin (World Resources Institute, Washington, DC); R. Lemmelä (National Board of Waters and the Environment, Helsinki, Finland); A. Leskelä (Finnish Committee on Large Dams, Vantaa); H. Löffler (Limnologisches Institut der Ö.A.W., Wien, Austria); J. Lundager Jensen (Det Danske Hedeselskab, Slagelse, Denmark); J. Maaranta (Imatran Voima O.Y., Vantaa, Finland); U. Mäkirinta (University of Oulu, Finland); A. Mallik and H. Rasiid (Lakehead University, Thunder Bay, Canada); N. Marjanovic (Belgrade, Serbia); R. R. Mason (U. S. Geological Survey, Raleigh, NC); P. J. McCurry (Environment Canada, Guelph); D. Mihail (Romanian National Committee on Large Dams, Bucharest); L. P. Mikhailov and V. D. Novozhenin (Hydroproject Institute, Moscow); J. Miller (Water Research Institute, Prague, Czech Republic); I. Obrusnik (Czech Hydrometeorological Institute, Prague); G. G. Patterson (U. S. Geological Survey, Columbia, SC); B. Pejaković (J. V. P. Hrvatska Vodoprivreda, Split, Croatia); G. E. Petts (University of Technology, Loughborough, UK); G. Piacentini (Autorità di Bacino del Fiume Po, Parma, Italy); R. Pol (Environment Canada, Moncton); D. Pontius (American Rivers, Washington, DC); A. Rodriguez (Texas Water

Development Board, Austin, TX); D. Rosenber (Freshwater Institute, Winnipeg, Canada); Ruhrverband, Hauptabteilung Talsperrenwesen (Esser Germany); R. D. Schopp (U.S. Geological Survey, West Trenton, NJ); E. Selvig (University of Oslo, Norway); P. G. Sly (Rawson Academy of Aquatic Sciences, Ottawa, Canada); P. Squires (Environment Canada, Yellowknife); State Committee for Hydrometeorology (Moscow); B. Statzner (Université Claude Bernard, Lyon, France); R. Sværre (Norges Vassdrags og Energiwerk, Narvik, Norway); R. Thepot (Etablissement Public d'Aménagement de la Loire et de ses Affluents, Orléans, France); H. Tómasson (National Energy Authority, Reykjavik, Iceland); L. Votruba (Prague, Czech Republic); T. Vučetić (Meteorological and Hydrometeorological Service, Zagreb, Croatia); V. S. Vugiński (State Hydrological Institute, St. Petersburg, Russia); Wasser- und Schifffahrtsverwaltung des Bundes (Aurich, Germany); K. Weckerle (München, Germany); W. A. R. Whitcomb (Manitoba Hydro, Winnipeg, Canada); D. D. Williams (University of Toronto, Canada); S. Winberg (Kristianstads Kommun, Kristianstad, Sweden); J. Yagüe (Comisión Internacional de Grandes Presas, Madrid, Spain); and M. J. Zimmerman (U.S. Geological Survey, Marlborough, MA). We also acknowledge S. Söderlind for drawing the maps; C. Andersson for assistance in compiling data for Norway, Sweden, Finland, and Denmark; and T. Elmquist, R. Jansson, M. E. Johansson, B. G. Jonsson, S. D. Wilson, and two anonymous reviewers for valuable comments on the manuscript.

AAAS–Newcomb Cleveland Prize

To Be Awarded for a Report, Research Article, or an Article Published in *Science*

The AAAS–Newcomb Cleveland Prize is awarded to the author of an outstanding paper published in *Science*. The value of the prize is \$5000; the winner also receives a bronze medal. The current competition period began with the 3 June 1994 issue and ends with the issue of 26 May 1995.

Reports, Research Articles, and Articles that include original research data, theories, or syntheses and are fundamental contributions to basic knowledge or technical achievements of far-reaching consequence are eligible for consideration for the prize. The paper must be a first-time publication of the author's own work. Reference to pertinent earlier work by the author may be included to give perspective.

Throughout the competition period, readers are

invited to nominate papers appearing in the Reports, Research Articles, or Articles sections. Nominations must be typed, and the following information provided: the title of the paper, issue in which it was published, author's name, and a brief statement of justification for nomination. Nominations should be submitted to the AAAS–Newcomb Cleveland Prize, AAAS, Room 924, 1333 H Street, NW, Washington, DC 20005, and **must be received on or before 30 June 1995**. Final selection will rest with a panel of distinguished scientists appointed by the editor-in-chief of *Science*.

The award will be presented at the 1996 AAAS annual meeting. In cases of multiple authorship, the prize will be divided equally between or among the authors.